

## Internship in theoretical nuclear physics

---

### Study of the fusion hindrance in reactions leading to the synthesis of super-heavy elements

---

One of the main questions in nuclear physics is why is there a finite number of nuclides, and, in particular, what are the heaviest possible nuclear objects in the region of super-heavy elements (SHE)? This limited number is related to their stability, as it is well known. Consequently, challenging research programmes are devoted to the study of very exotic nuclide at the limit of stability where new properties may appear and where one might have direct access to the specific features of nuclear matter being responsible for their existence. These limits are located at the edges the chart Segrè in regions extreme proton-neutron ratios (proton/neutron driplines) and at highest masses  $A$  and atomic numbers  $Z$ .

The so-called super-heavy nuclei (SHN) are expected to exist beyond the liquid drop limit of existence defined by a vanishing fission barrier thanks to the quantum mechanical shell effects. These nuclei are particularly interesting because they are at the limit between few body and large  $n$ -body physics: magic proton and neutron numbers,  $Z$  and  $N$ , are replaced by a magic region or island extended in  $Z$  and  $N$ .

Synthesis of these very and super-heavy nuclei by fusion-evaporation reactions is an experimental challenge due to the extremely low cross-sections. Modelling the complete reaction in order to guide the experiments is also a difficult challenge because models developed for lighter nuclei cannot be simply extrapolated. Fusion reactions are hindered with respect to what is observed with light nuclei because of the very strong Coulomb interaction being enhanced by the strong repulsion created by the large number of positive charges (protons) in the system. Due to its competition with the attractive strong (nuclear) force in a highly dynamic regime. The predictive power of the models is low although the origin of the hindrance phenomenon is qualitatively well understood. Quantitative ambiguities are large enough to observe few orders of magnitude differences in the fusion probabilities calculated by different models. A small change of the cross-section could result in many months being required to successfully perform experiments.

In such a context, the main purpose of the theoretical researches performed at GANIL on the synthesis of SHE is aimed to find ways to assess the models in order to improve their predictive power.

During this internship, the student will study a new mechanism that could significantly contribute to the fusion hindrance. The purpose of the study is to link a theoretical model with a phenomenological approach to see if the former can explain the latter.

**Expected skills:** Nuclear physics, modelling

This work can be pursued by the PhD-thesis entitled *Study of reaction mechanisms for the synthesis of super-heavy elements*.

**Contact:**

David Boilley  
GANIL, BP 55027, 14076 Caen France  
Tel: +33 (0)2 31 45 47 81  
Mail: [boilley@ganil.fr](mailto:boilley@ganil.fr)